

204070-2722001




FIG. 1

2/14

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted February 14, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.





FIG. 2

3/14



Excimer configuration
of Pyrene pairs

FIG. 3

201010 212600T




FIG. 4

4/14

5/14




FIG. 5

6/14

Addition of 0.48 μ M Zip3R to
0.08 μ M Zip3F




FIG. 6

7/14




FIG. 7

8/14

20000-22000





FIG. 8


9/14



- 1 - Autocamtide II phosphorylated by CaMKII in the presence of Ca^{2+} and Calmodulin
- 2 - Zip4S phosphorylated by PKA
- 3 - Zip4S phosphorylated by CaMKII in the presence of Ca^{2+} and Calmodulin
- 4 - Autocamtide II phosphorylated by CaMKII in the presence of Calmodulin (Ca^{2+} absent)

FIG. 9


10/14



1 - zip4S
2 - zip4T

FIG. 10

11/14



1 - zip4S
2 - zip4SP

FIG. 11

12/14

204010-2F2E00F





FIG. 12

13/14



- 1 - 0.08 μM Zip4SF
- 2 - 0.08 μM Zip4SF + 0.48 μM Zip4SR
- 3 - 0.08 μM Zip4SF + 0.48 μM Zip4SR + 0.03 μM PKA

FIG. 13




FIG. 14